



2.0
ANCC
CONTACT HOURS

How to recognize and respond to refeeding

To avoid potentially lethal consequences, assess a malnourished patient carefully before restarting nutrition by any route.

By Mary Ann Yantis, RN, PhD, and
Robyn Velandar, RD, LD, CNSD

You're caring for Betty Dawson, 73, who's newly admitted with dehydration and malnutrition. She's 5 feet 6 inches tall and weighs 121 pounds (55 kg). Her admission basic metabolic panel is normal except for a slight increase in serum sodium (149 mEq/L; normal range, 136 to 145 mEq/L) and a decrease in potassium (3.3 mEq/L; normal range, 3.5 to 5.2 mEq/L). Her baseline magnesium level is 2.2 mg/dL (normal range, 1.8 to 2.6 mg/dL), and her baseline phosphorus level is 3 mg/dL (normal range, 2.5 to 4.5 mg/dL). She's initially treated with dextrose 5% in 0.45% sodium chloride solution for hydration and I.V. potassium to correct her hypokalemia. Ms. Dawson is diagnosed with inflammatory bowel disease and malnutrition.

On hospital day 2, she's started on parenteral nutrition (PN) at a rate that delivers 25 to 35 kcal/kg or 1,375 to 1,925 kcal/day, and her condition is stable. But when you assess her early in the morning on hospital day 3, you note lethargy, shortness of breath, and muscle weakness. All of her vital signs are normal except her pulse, which has become irregular. Would you recognize that these nonspecific signs might indicate a potentially life-threatening disorder?

Severely malnourished patients can experience significant fluid shifts and electrolyte imbalances after aggressive nutritional support is initiated. This potentially lethal disorder, known as refeeding syndrome, usually is associated with PN, but it also can occur with enteral nutrition, oral intake, or dextrose-containing I.V. fluids.¹

The background is an abstract composition of soft, blended colors. A gradient transitions from a deep purple at the top to a warm orange at the bottom. A sharp, diagonal line, appearing as a reflection or a boundary, runs from the top-left towards the bottom-right, adding a sense of movement and depth to the design.

syndrome

Who's at risk for malnutrition?^{1,3}

Assess and closely monitor patients with these conditions:

- chronic alcoholism
- anorexia nervosa
- cancer, especially when on chemotherapy
- massive weight loss after being morbidly obese
- chronic gastrointestinal disease, such as Crohn's disease
- prolonged I.V. hydration with nonnutritional fluids
- stroke
- protein-calorie malnutrition
- prolonged fasting, defined as no oral intake for more than 7 days
- loss of 10% of body weight or more within the preceding 4 weeks
- decreased intake for any reason after surgery.

Although information about refeeding syndrome is available in nutrition journals, it isn't widely discussed in nursing literature. We'll fill that gap with this article by discussing how to identify this dangerous complication and what to do about it.

Disorder discovered

This syndrome was first observed and described after World War II when victims of starvation were noted to experience cardiac or neurologic dysfunction or both after being reintroduced to food. Today, we rarely see patients who are as severely malnourished as these people were, but protein-calorie malnutrition is common among hospitalized patients: Almost one-third are malnourished and nearly two-thirds are at risk for malnutrition.²

Electrolyte disturbances (primarily decreased levels of phosphorus, magnesium, or potassium) occur immediately upon the rapid initiation of refeeding—commonly within 12 or 72 hours—and can continue for the next 2 to 7 days. Cardiac complications can develop within the first week, often within the first 24 to 48 hours, with neuro-

logic signs and symptoms developing somewhat later.³

The following patients are at particular risk for refeeding syndrome:

Hospitalized patients.

Refeeding syndrome occurs primarily in hospitals. One study found that in 51 patients who received vigorous nutritional support, 80% of the patients developed depletions in potassium, magnesium, or phosphorus. Ninety-three percent of the patients in this study who were considered "at nutrition risk" and 74% of those "not at nutrition risk" had electrolyte imbalances.⁴ These results show the importance of closely monitoring electrolytes in all patients who are receiving enteral or PN support.

Children. Although refeeding syndrome is reported most often in adults, it also can occur in children. In a review of the literature, we found reports of 27 children who developed this syndrome after initiation of oral or enteral feeding; nine of them died as a result of related complications.⁵

Patients with chronic malnutrition. Refeeding syndrome can develop in patients who are chronically malnourished for any

reason (see *Who's at risk for malnutrition?*). At particular risk are those who:

- have lost at least 10% of their weight within the preceding 4 weeks
- haven't eaten for more than 7 days, such as a patient experiencing nausea and vomiting or anorexia from chemotherapy
- have chronic alcoholism
- have had rapid and massive weight loss after previously being obese.⁶

Understanding starvation

To understand what happens during refeeding syndrome, first review the pathophysiology of malnutrition. Normally, glucose is the body's preferred fuel, coming from the intake of carbohydrates. As the malnourished body loses access to carbohydrates, it shifts to catabolism of fat and protein. With this shift, the body's production of insulin drops in response to a reduced availability of glucose. This adaptive change to protein breakdown during prolonged malnutrition also leads to a gradual loss of cellular and muscle mass, often resulting in atrophy of vital organs and other internal structures, including the heart, lungs, liver, and intestines.

Serious complications may occur as respiratory and cardiac function declines due to muscular wasting and fluid and electrolyte imbalances. Metabolic rate, cardiac output, hemoglobin levels, and renal concentration capacity also decrease.⁷ The body is now surviving by very slowly consuming itself.


As you think of these changes, you probably envision a thin, emaciated patient, but even an

How electrolyte levels change in refeeding syndrome

Here's what happens once a malnourished patient begins aggressive nutritional support:


Step 1

Glucose (G) enters bloodstream by enteral, parenteral, oral, or I.V. route.




Step 2

Pancreas secretes insulin (I) in response to increase in serum glucose.



Step 3

Insulin promotes cellular uptake of glucose, potassium (K⁺), magnesium (Mg⁺⁺), and phosphorus (P⁻). Low serum electrolyte levels for potassium, phosphorus, and magnesium result.



overweight or obese patient can be malnourished.

What goes wrong

When a malnourished patient is given aggressive nutritional sup-

port, such as PN, a number of events ensue. These are primarily driven by the change in insulin secretion as a result of the shift from protein metabolism to carbohydrate metabolism. The

increase in glucose levels, which results from the composition of the nutritional support formula, increases insulin release by the pancreas. This in turn promotes cellular uptake of glucose

Watch for these signs and symptoms³

Electrolyte imbalance

- hypokalemia
- hypophosphatemia
- hypomagnesemia

Neurologic

- weakness
- seizure
- paresthesia
- altered mental status
- paralysis
- tetany

Musculoskeletal

- weakness
- myalgia

Hematologic

- thrombocytopenia
- platelet dysfunction
- anemia
- infections
- bleeding

Respiratory

- hypoxia

Gastrointestinal

- anorexia
- abdominal pain
- constipation
- diarrhea

Cardiovascular

- dysrhythmias
- hypotension

Renal

- edema
- elevated blood urea nitrogen
- elevated creatinine

Metabolic

- metabolic alkalosis
- metabolic acidosis

Nutritional

- vitamin deficiency, such as deficiency of vitamin B₁ (thiamine)
- low albumin and prealbumin levels.

along with electrolytes, primarily phosphorus, magnesium, and potassium. The result can be a life-threatening depletion of these vital electrolytes.^{1,8} (See *How electrolyte levels change in refeeding syndrome*.)

First, let's look at some normal functions of phosphate. It's needed to produce adenosine triphosphate (ATP), which provides energy for almost all cellular functions. Phosphate is an essential part of RNA and DNA, and it's needed in red blood cells for 2,3-diphosphoglycerate production for easier release of oxygen to the tissues. Patients with signs and symptoms of hypophosphatemia or phosphate levels below 2 mg/dL require oral or I.V. phosphate replacement.⁹

Refeeding-induced severe hypophosphatemia (serum concentration less than 1 mg/dL) can result in respiratory failure from a decrease in available ATP, which is needed to maintain the diaphragm's normal contractility. In addition, hypophosphatemia can cause red and white blood cell dysfunction, muscle weakness, and seizures. Other factors that can contribute to hypophosphatemia include vitamin D deficiency and excessive intake of antacids, which block phosphate absorption.¹⁰

Hypokalemia (serum levels below 3.5 mEq/L) and hypomagnesemia (serum levels below 1.8 mg/dL) are also frequently associated with refeeding syndrome. Mild decreases of potassium and magnesium may cause nausea, vomiting, constipation, diarrhea, muscle twitching, or weakness. A more severe depletion of the serum concentrations of potassium and magnesium can cause

dysrhythmias, cardiac dysfunction, skeletal muscle weakness, seizures, and metabolic acidosis.¹⁰

Collaboration is key

To prevent refeeding syndrome or identify early signs and symptoms, you'll need to communicate and coordinate often with healthcare providers, dietitians, pharmacists, and other nurses. (See *Watch for these signs and symptoms*.) Your role centers on identifying patients who are at risk and recognizing signs and symptoms early on. You may be the first person to take a patient's history and perform a physical assessment, so you may be the first to learn about her preadmission weight loss or problems that prevent adequate nutritional intake, such as nausea, vomiting, and dysphagia. Once you identify risk factors for refeeding syndrome, alert her healthcare provider and a registered dietitian.

Enter the dietitian

The dietitian can identify the level of malnutrition (mild, moderate, or severe) based on the patient's rate of weight loss and her current body weight, and by evaluating the patient's visceral protein status based on levels of serum albumin, prealbumin, and transferrin or her total lymphocyte count. A dietitian will help determine the appropriate methods of nutrition support (parenteral, enteral, or oral) and request additional lab tests.

Tailoring nutrition support to the patient's condition requires the calculation of her basic energy requirement, or basal energy expenditure—the amount of energy required to maintain the

body's normal metabolic activity at rest. The dietitian is well educated in evaluating a patient's caloric need and can provide guidance on how and when to increase the nutritional load of the nutrition support formula.


Your role in patient care

If your patient is at risk for refeeding syndrome, follow these guidelines to prevent or minimize problems.

- *Monitor serum electrolyte levels and make sure they're normal before beginning nutritional support.* Watch particularly for abnormal levels of potassium, phosphate, and magnesium. If they aren't normal, they should be corrected promptly. These levels also need to be checked every 6 hours, 12 hours, or daily for at least 3 days after nutrition is initiated.¹

- *Carefully assess BP, pulse rate, and intake and output.* Your patient may be dehydrated and hypovolemic before she begins therapy. Carefully restore volume and sodium if necessary. If your patient is severely malnourished, her fluid intake may need to be restricted. She also should avoid excessive sodium intake. She may have an altered cardiac rate or rhythm at the start of therapy, and changes in pulse rates can indicate a change in her clinical status. Tell her healthcare provider about any such clinical changes.

- *Closely monitor the refeeding rate.* When resuming nutrition, start slowly (especially when PN is used), at a rate that delivers 15 to 20 kcal/kg/day (about 1,000 kcal/day for adults) for the first 1 to 3 days, before gradually advancing to the desired daily



During refeeding, a malnourished patient can rapidly develop fluid overload, so be sure to monitor her for abnormal lung sounds.

levels over 5 to 7 days. The healthcare provider will advance nutritional support when the serum electrolytes are close to the normal range or are actively being replaced.

- *Provide patient education.* If a patient is showing signs of refeeding syndrome after oral food intake, you should help slow down the reintroduction of carbohydrates by encouraging her to eat small meals. Tell her to choose supplements or snacks that are low in sugar (for example, sugar-free juices or sugar-free puddings or gelatins) and fluids that don't contain carbohydrates. Eating food that's high in potassium can help reverse potassium repletion but at a much slower rate than when potassium is given I.V. High-potassium foods include melons, milk, bananas, apricots, oranges, and potatoes. Giving carbohydrate-rich snacks along with a high-protein source (for example, crackers with cheese and peanut butter) can help slow the rate of digestion and may slow the rate of insulin secretion.

- *Monitor phosphorus, magnesium, and potassium levels daily for the first week or until levels become stable.* Life-threatening changes in these electrolytes commonly occur in the first 1 to 3 days of therapy. Notify the primary care provider if abnormalities develop in any of these values. You should continue closely observing your patient during correction of any of these electrolyte abnormalities. For example, while phosphorus is being administered, monitor your patient for possible adverse reactions such as tetany, hypotension, hyperkalemia, and hypernatremia. Be aware of medications that may influence the development of hypophosphatemia, hypokalemia, or hypomagnesemia. Consult with the pharmacist if necessary to identify additional medications that may contribute to electrolyte imbalances.

- *Assess vitamin levels.* Malnourished patients, especially those with alcoholism, are likely to have deficiencies of vitamins, including thiamine. Thiamine deficiency can result in confusion, ataxia, and coma. As a general rule, any malnourished patient at risk for refeeding syndrome should receive thiamine supplementation.

- *Meticulously document fluid intake and output and weight.* During refeeding, a malnourished patient can rapidly develop fluid overload, so be sure to monitor her for abnormal lung sounds, which can be the first sign of fluid overload. Also monitor her intake and output and weigh her daily. Assess for edema of her lower extremities, which may be an early sign of fluid overload. Weight gain of more than 0.5 pound (0.23 kg) a day or

3.3 pounds (1.5 kg) a week probably represents fluid retention. If excessive weight gain occurs, notify the healthcare provider.

- **Monitor blood glucose levels frequently.** Your patient can rapidly become hyperglycemic when she's receiving glucose replacement. Generally, during the initial refeeding stage, glucose levels should be kept at or below 200 mg/dL. Closely monitor your patient for additional sources of glucose she's receiving; for example, dextrose in I.V. fluids or drugs that can affect serum glucose such as corticosteroids and beta-blockers.

- **Monitor for neurologic signs and symptoms.** Your patient with refeeding syndrome may develop muscle weakness, tremors, paresthesias, and seizures. Institute seizure precautions. In addition, she may have cognitive changes,

including irritability and confusion. These cognitive changes make her safety a priority. Institute fall risk procedures.

Remain vigilant

Although not all patients who receive nutritional support develop refeeding syndrome, you must be aware of this condition and identify and closely monitor those patients at risk. Your vigilance can help protect patients from the potentially devastating consequences of refeeding syndrome. ♦

REFERENCES

1. McCray S, Walker S, Parrish CR. Much ado about refeeding. *Pract Gastroenterol.* 2005;30(1):26-44.
2. Thomas DR, Zdrowski CD, Wilson MM, et al. Malnutrition in subacute care. *Am J Clin Nutr.* 2002;75(2):308-313.
3. Crook MA, Hally V, Panteli JV. The importance of refeeding syndrome. *Nutrition.* 2001;17(7-8):632-637.
4. Flesher ME, Archer KA, Leslie BD, et al. Assessing the metabolic and clinical consequences of early enteral feeding in the malnourished patient. *JPEN J Parenter Enteral Nutr.* 2005;29(2):108-117.
5. Afzal NA, Addai S, Fagbemi A, et al. Refeeding syndrome with enteral nutrition in children: a case report, literature review and clinical guidelines. *Clin Nutr.* 2002;21(6):515-520.
6. Lauts NM. Management of the patient with refeeding syndrome. *J Infus Nurs.* 2005;28(5):337-342.
7. Ladadge E. Refeeding syndrome. *ORL Head Neck Nurs.* 2003;21(3):18-20.
8. Worthington PH. *Practical Aspects of Nutrition Support: An Advanced Practice Guide.* Philadelphia, PA: Saunders; 2004.
9. Mailhot T, Richard AJ. Hypophosphatemia. <http://www.emedicine.com/emerg/topic278.htm>.
10. Kraft MD, Btaiche IF, Sacks GS. Review of refeeding syndrome. *Nutr Clin Pract.* 2005;20(6):625-633.

Mary Ann Yantis is an associate professor of nursing at the Louise Herrington School of Nursing at Baylor University in Dallas, Tex., and her daughter Robyn Velandier is a clinical dietitian at Adventist Medical Center in Portland, Ore.

The authors have disclosed that they have no financial relationships related to this article.

Adapted from Yantis MA, Velandier R. How to recognize and respond to refeeding syndrome. *Nursing.* 2008;38(5):34-40.

For more than 20 additional continuing education articles related to nutrition, go to NursingCenter.com/CE.

CE CONNECTION

Earn CE credit online:

Go to <http://www.nursingcenter.com/NCC> and receive a certificate within minutes.

INSTRUCTIONS

How to recognize and respond to refeeding syndrome

TEST INSTRUCTIONS

- To take the test online, go to our secure Web site at <http://www.nursingcenter.com/NCC>.
- On the print form, record your answers in the test answer section of the CE enrollment form on page 21. Each question has only one correct answer. You may make copies of these forms.
- Complete the registration information and course evaluation. Mail the completed form and registration fee of \$21.95 to: Lippincott Williams & Wilkins, CE Group, 2710 Yorktowne Blvd., Brick, NJ 08723. We will mail your certificate in 4 to 6 weeks. For faster service, include a fax number and we will fax your certificate within 2 business days of receiving your enrollment form.
- You will receive your CE certificate of earned contact hours and an answer key to review your results. There is no minimum passing grade.
- Registration deadline is June 30, 2011.

DISCOUNTS and CUSTOMER SERVICE

- Send two or more tests in any nursing journal published by Lippincott Williams & Wilkins together and deduct \$0.95 from the price of each test.
- We also offer CE accounts for hospitals and other health care facilities on nursingcenter.com. Call 1-800-787-8985 for details.

PROVIDER ACCREDITATION

Lippincott Williams & Wilkins, publisher of *Nursing2009 Critical Care*, will award 2.0 contact hours for this continuing nursing education activity.

Lippincott Williams & Wilkins is accredited as a provider of continuing nursing education by the American Nurses Credentialing Center's Commission on Accreditation.

This activity is also provider approved by the California Board of Registered Nursing, Provider Number CEP 11749 for 2.0 contact hours. Lippincott Williams & Wilkins is also an approved provider of continuing nursing education by the District of Columbia and Florida #FBN2454. LWW home study activities are classified for Texas nursing continuing education requirements as Type I.

Your certificate is valid in all states.